

You are receiving this package because you or your child are registered for the Scuba Certification Adventure at Sea Base for the 2026 season. This will be a fun and exciting adventure for all attending, but there is some preliminary work that needs to be completed prior to your arrival at Sea Base. This package contains information about the adventure medical requirements, medical forms and releases. This year we are using CampDoc for our Medical documents collection system. Your Crew Leader will supply CampDoc with the names and information needed to set up your CampDoc profile. You will then be able to complete and upload the forms on the CampDoc system. Some forms are completed on the system and some need to be uploaded. The Scouting America Part C and the RSTC Diver Medical Questionnaire page 3 will need to be completed by the physician and uploaded to the system.

### All Medical forms and additional documents are due by the following dates:

- Spring Crews January 1, 2026
- Summer Crews- March 1, 2026

Documents returned after the deadline may affect your crew's ability to participate in their full adventure.

This package contains the following:

- Sea Base Participant Health & Safety Advisory- This is information regarding Scouting America Medical Requirements for Scuba Diving.
- Scouting America Annual Health and Medical Record (AMHR)-All sections must be completed in their entirety.
   Part C must be completed and signed by a physician (MD or DO only). Parts A and B are completed on the
   CampDoc system. Part C will need to be uploaded
- Recreational Scuba Training Council (RSTC) Diver Medical Participant Questionnaire-The Physicians Evaluation
   Form must be signed by a physician (MD or DO only) even if all questions on pages 1 and 2 were answered "NO".

   Pages 1 and 2 are completed on the CampDoc system. Page 3 needs to be uploaded.
- Florida Addendum Notice to the Minor Child's Parent Completed on the CampDoc system.
- **PADI Release for General Training Form-** The Non-Agency Disclosure & Liability Release Agreements must be signed by both the youth, even if under 18 years of age, and parents. Completed on the CampDoc system.
- PADI Safe Diving Practices Form-Must be signed by both the youth, even if under 18 years of age, and parents. Completed on the CampDoc system.
- \*Sea Base Dietary Notification form-only if food allergies present, this form is available on our website.

Additional supporting documentation that must be submitted in addition to the above documents:

- Participants over 18 years of age must submit proof of the following Scouting America Adult Trainings:
  - Safeguarding Youth
  - Safety Afloat
  - Safe Swim Defense
  - Weather Hazards
- Scouting America Membership card Presented at time of arrival for both youth and adult.
- Copy of front and back of Certification Card (Scuba Cert. Rescue Option Only)

We advise all participants and parents to read the <u>Participant Guide</u> and <u>PowerPoint</u> for your adventure. Both items are found on the Sea Base website.

Feel free to contact the Sea Base Scuba Department if you have any questions relating to this information.

# Sea Base Participant Health & Safety Advisory

This document is meant to serve as a guide to understand health and safety considerations for all Sea Base Program in the Florida Keys, USVI and the Bahamas. This document also outlines contradictions that may preclude participation in some of our programs/ activities per Scouting America Policy. Please note that Sea Base does not dictate policies set by Scouting America in regards to Scuba Diving and Snorkeling. Sea Base, Scouting America is an industry leader in maritime adventures with an excellent safety record. If you have questions regarding medical policies, medical concerns, or medical approval, please contact Sea Base at 305-664-4173.

### General Health & Safety Info

Sea Base Experience: Sea Base adventures are not risk-free. Participants must follow safety measures and take personal responsibility for their health and safety. Climate conditions include temperatures ranging from 45 to 95 degrees, high humidity, heat indexes reaching 110 degrees, and frequent, sometimes severe tropical weather. Prolonged, rigorous activities include snorkeling, scuba diving, kayaking, canoeing, sailing, hiking, and others. It is the responsibility of participants, participant parents or guardians, participant health-care teams, and unit leaders to see that each individual—youth or adult—can safely take part in Sea Base adventures.

Adult Participants: It is the role of accompanying adults to ensure youth safety. Because of this, adult participants must arrive in good physical condition and have no medical conditions that could result in diverting the Sea Base staff's attention away from youth participants.

Participant Ability Level: Sea Base Participants must be able to:

- Swim in a strong manner
- Climb a 6-foot ladder, unassisted, in inclement weather, from the water onto a rocking vessel
- Self-rescue if found overboard in inclement weather

**Trained Leadership:** Each crew is required to have at least one adult who is trained in wilderness first aid and CPR or has a greater professional medical certification. This leader acts as the primary first response until emergency services arrives. There are no on-site facilities for treatment or extended care at Sea Base. Sea Base does not staff professional medical personnel.

Location: Sea Base adventures are conducted at sea, often far from land, with limited access to emergency services. Response times can be affected by weather, seas, and location, and can be delayed for hours. Individuals with medical conditions that require immediate or nearly immediate access to professional medical care should not attend Sea Base.

Right to Refuse: Sea Base reserves the right to deny participation based on health and safety concerns and/or medical history.

Special Needs or Medical Concerns: Any individual with special needs or medical concerns must have an onsite advocate who understands the individual's condition and treatment and who is prepared to provide support to the individual. Special Needs youth who do not meet certain attendance requirements may be eligible to attend COASTS designated programs.

Medications: Individuals requiring medication should continue medications as prescribed and bring an appropriate supply. Each crew must develop a plan to secure, lock, and dispense medication.

Scuba Participants: Persons with conditions listed as severe by the Recreational Scuba Training Council (RSTC) will not be permitted to scuba dive. Persons with conditions prohibited by Scouting America scuba policy will not be permitted to scuba dive. Various risk factors may exclude a person from scuba diving, either temporarily or permanently. Risk factors include, but are not limited to, ear and sinus problems, recent surgery, spontaneous pneumothorax, asthma or reactive airway disease, seizure disorders, diabetes, leukemia, sickle-cell disease, pregnancy, panic disorders, active psychosis, certain medications, and narcolepsy. For more specifics please see the <u>Absolute Medical Contradictions for SCUBA Section here.</u>

### **General Health Considerations**

Weight Limits: Those persons weighing more that 295 pounds will not be permitted to attend.

If a participant weighs more than 295 pounds, they will be sent home at their own expense. There will be weigh-ins upon arrival at Sea Base. There are no exceptions or waivers to the 295-pound weight limit

Allergies: Participants with allergies that may result in severe reactions or anaphylaxis should bring an adequate supply of epinephrine auto-injectors (EpiPen) to last up to three hours.

Sleep Apnea: Prospective participants with sleep apnea may participate at Sea Base if they have been medically cleared for participation, they understand that they are responsible to provide battery support for their CPAP and may not have access to electricity if participating in Bahamas, St. Thomas, St. Croix, Florida Keys Sailing, SCUBA Live Aboard, Keys Adventure or Out island programs.

### Recommendations Regarding Chronic Illness and/or Compromised Immune

**System:** Persons with chronic conditions and/or compromised immune systems should seek medical advice and education regarding medical risks associated with harsh marine environments before

participating. Individuals with open wounds or who are at risk for chronic illnesses or immune disorders should not attend Sea Base.

Hypertension (High Blood Pressure): Participants should have a blood pressure less than 140/90. Individuals with hypertension should have the condition treated and well-controlled before attending.

Recent Musculoskeletal Injuries and Orthopedic Surgery: Persons with musculoskeletal problems or orthopedic surgeries within the last six months must provide a letter from their treating physician to participate.

Psychological and Emotional Difficulties: Any condition should be well-controlled without the services of a mental health practitioner. Participants requiring medication must bring an ample supply and take as prescribed for the duration of their trip. If you are a SCUBA Participant please see the Absolute Medical Contradictions For Scuba Diving- Psychological & Emotional Difficulties Section.

Insulin-Dependent Diabetes Mellitus: Diabetes must be well-controlled. Hypoglycemia can lead to unconsciousness and drowning. <u>If you are a Scuba Participant</u>, please see the <u>Absolute Medical Contradictions For Scuba Diving-Insulin Dependent Diabetes Mellitus Section</u>.

Asthma & Reactive Airway Disease: Asthma must be well-controlled. Persons requiring use of medication and/or an inhaler must bring an ample supply. If you are a Scuba Participant, please see the Absolute Medical Contradictions for SCUBA Diving- Asthma & Reactive Airway Disease Section.

### Absolute Medical Contradictions for Snorkeling & Scuba Diving

Seizures (Epilepsy): Seizures while snorkeling or scuba diving are extremely dangerous and often fatal.

- Snorkeling or scuba diving as part of an official scouting activity is prohibited for participants with a history of seizures.
- No participant with a history of seizures or taking anti-epileptic medication may snorkel or scuba
  dive. Participants meeting these criteria must be wearing a USCG approved lifejacket and may not
  be wearing a mask anytime they are in the water. Participants meeting these criteria are allowed to
  swim in the water but must be accompanied by their buddy AND an adult who is trained to
  recognize symptoms of a seizure.
  - Prospective participants with a history of infant febrile seizures may be considered for snorkeling or scuba diving after formal consultation with a neurologist.

### Absolute Medical Contradictions for Scuba

### Absolute medical contradictions for scuba diving:

- Asthma
- Epilepsy / Seizures
- Insulin dependent diabetes
- Anxiety requiring medication
- Narcolepsy
- Spontaneous pneumothorax

- Exceeding the weight limit of 295 pounds
- Multiple (more than one) medication for ADD, ADHD or depression
- Blood thinners
- Migraines with auras
- Hearing loss in one ear

### Risk factors include, but are not limited to:

- Ear and sinus problems
- Sickle-cell disease
- Recent surgery
- Pregnancy
- Chemotherapy
- · Panic disorders

- Leukemia
- Active psychosis
- Certain medications
- Migraines requiring medication
- Blood thinning medication
- Pacemakers

NOTE: For additional information read the detailed description on the following pages. Not every disqualifying medical condition for Scouting America scuba diving is listed. If you require further clarification regarding an unlisted medical condition and scuba diving with the Scouting America, please contact the Sea Base Scuba Department.

### Insulin-Dependent Diabetes Mellitus:

Diabetes must be well-controlled. Hypoglycemia can lead to unconsciousness and drowning.

- Diving as part of an official scouting activity is prohibited for the following:
  - o For persons under age 18 with the diagnosis of diabetes.
  - Persons using insulin to control diabetes.
  - Persons with diabetes, who are non-insulin dependent and who have had recurrent problems and/or hospitalizations for diabetic problems.
  - o Persons with any HbA1c test greater than 7.0 in the previous 12 months.
  - Persons having a documented or suspected hypoglycemic event requiring treatment or assessment in the previous 12 months.
- Diabetes is considered well-controlled when the following are met:

 The acceptable oral medications for diabetic control are as single agents only: metformin and metformin analogs; DPP-4 inhibitors (sitagliptin, vildagliptin, alogliptin, saxagliptin and linagliptin); or SGLT2 inhibitors and analogies.

Persons who control their diabetes with exercise and diet (without the aid of medication, except metformin) and document HbA1c test value less than 7.0 in the last 6 months) may be approved to scuba dive.

Participants that are insulin dependent, youth or adult, will not be cleared to scuba dive.

Those that are currently scuba certified or have a physician's approval for scuba diving will not be cleared to scuba dive at Sea Base. Participants younger than 18 years of age with diabetes will not be cleared to scuba dive. There are no exceptions, exclusions or waivers to this policy.

### Asthma & Reactive Airway Disease:

Diving as part of an official scouting activity is prohibited for persons being treated for asthma or reactive airway disease.

- Persons with a history of asthma who have been asymptomatic and have not used medications to control asthma for five years or more may be allowed to scuba dive if resolution of asthma is specifically confirmed by their physician and includes provocative pulmonary function testing conducted by a pulmonologist.
  - o Provocative testing can include exercise, hypertonic saline, a hyperpnea test, etc.

Participants, youth or adult, with asthma will not be cleared to scuba dive. The predisposing factors, severity of attacks or intermittent asthma does not change this Scouting America policy. Those that are currently scuba certified or have a physician's approval for scuba diving will not be cleared to scuba dive at Sea Base. There are no exceptions, exclusions or waivers to this policy.

### Psychological and Emotional Difficulties (ADD, ADHD, anxiety and depression):

Any condition should be well controlled.

- Diving as part of an official scouting activity is prohibited for:
  - o Participants taking more than one medication for any of these conditions.
  - o Participants with anxiety disorder requiring any medication.

Several medications are <u>NOT COMPATIBLE</u> with the hyperbaric stresses of scuba diving. All medications MUST be listed on the Scouting America Annual Health and Medical Record. Those participants on multiple medications (more than one) for ADD, ADHD, Depression or any psychological condition <u>will not be cleared for scuba diving</u>. Those participants requiring medication to control Anxiety will not be cleared for scuba diving. There are no exceptions, exclusions or waivers to this policy.

### **Concussions:**

Those participants who have suffered a concussion and any side effects from the concussion should contact DAN and Sea Base to discuss this issue. Please provide the following information on the Scouting America Medical:

- 1. When did the injury occur?
- 2. Was there any loss of consciousness, inability to recollect events?
- 3. Were there reports of disorientation after the incident?
- 4. Have there been any lingering affects?

### Closing:

Please note that the final decision for participation in all Sea Base programs is at the discretion of the Sea Base Medical Director. The decision is final and we are unable to reverse or alter that decision. No waivers will be issued.



### Part A: Informed Consent, Release Agreement, and Authorization

	<u> </u>
Full name:	High-adventure base participants:
Date of birth:	Expedition/crew No.:
Date of Shan	or staff position:
Informed Consent, Release Agreement, and Authorization	
I understand that participation in Scouting activities involves the risk of personal injury, including death, due to the physical, mental, and emotional challenges in the activities offered. Information about those activities may be obtained from the venue, activity coordinators, or your local council. I also understand that participation in these activities is entirely voluntary and requires participants to follow instructions and abide by all applicable rules and the standards of conduct.  In case of an emergency involving me or my child, I understand that efforts will be made to contact the individual listed as the emergency contact person by the medical provider and/or adult leader. In the event that this person cannot be reached, permission is hereby given to the medical provider selected by the adult leader in charge to secure proper treatment, including hospitalization, anesthesia, surgery, or injections of medication for me or my child. Medical providers are authorized to disclose protected health information to the adult in charge, camp medical staff, camp management, and/or any physician or health-care provider involved in providing medical care to the participant. Protected Health Information/Confidential Health Information (PHI/CHI) under the Standards for Privacy of Individually Identifiable Health Information, 45 C.F.R. §§160.103, 164.501, etc. seq., as amended from time to time, includes examination findings, test results, and treatment provided for purposes of medical evaluation of the participant, follow-up and communication with the participant's parents or guardian, and/or determination of the participant's ability to continue in the program. I further authorize the sharing of the information on this form with any BSA volunteers or professionals who need to know of medical conditions that may require special consideration in conducting Scouting activities.  With appreciation of the dangers and risks associated with programs and activities, on my	lalso hereby assign and grant to the local council and the Boy Scouts of America, as well as their authorized representatives, the right and permission to use and publish the photographs/film/videotapes/electronic representations and/or sound recordings made of me or my child at all Scouting activities, and I hereby release the Boy Scouts of America, the local council, the activity coordinators, and all employees, volunteers, related parties, or other organizations associated with the activity from any and all liability fromsuch use and publication. I further authorize the reproduction, sale, copyright, exhibit, broadcast, electronic storage, and/or distribution of said photographs/film/videotapes/electronic representations and/or sound recordings without limitation at the discretion of the BSA, and I specifically waive any right to any compensation I may have for any of the foregoing.  Every person who furnishes any BB device to any minor, without the express or implied permission of the parent or legal guardian of the minor, is guilty of a misdemeanor. (California Penal Code Section 19915[a]) My signature below on this form indicates my permission.  I give permission for my child to use a BB device. (Note: Not all events will include BB devices.)  Checking this box indicates you DO NOT want your child to use a BB device.  NOTE: Due to the nature of programs and activities, the Boy Scouts of America and local councils cannot continually monitor compliance of program participants or any limitations imposed upon them by parents or medical providers. However, so that leaders can be as familiar as possible with any limitations, list any restrictions imposed on a child participant in connection with programs or activities below.
own behalf and/or on behalf of my child, I hereby fully and completely release and waive any and all claims for personal injury, death, or loss that may arise against the Boy Scouts	
of America, the local council, the activity coordinators, and all employees, volunteers, related parties, or other organizations associated with any program or activity.	List participant restrictions, if any: □ None
related parties, or other organizations associated with any program or activity.	
Iunderstand that, if any information I/we have provided is found to be inaccurate, it may limit and/o Philmont Scout Ranch, Philmont Training Center, Northern Tier, Sea Base, or the Summit Bechtel Reand weight requirements and restrictions, and understand that the participant will not be allowed to the participant has permission to engage in all high-adventure activities described, except as parent or guardian's signature is required.	serve, I have also read and understand the supplemental risk advisories, including height owed to participate in applicable high-adventure programs if those requirements are not
Participant's signature:	Date:
Parent/guardian signature for youth:(If participantis und	Date:
приничний	
Complete this section for youth participants only:	
Adults Authorized to Take Youth to and From Events:	
You must designate at least one adult. Please include a phone number.	
Name:	Name:
Phone:	Phone:
Adults NOT Authorized to Take Youth to and From Events:	
Addition Addition Legito Take Touth to and Front Events.	



Part B1: General Information/Health History



Full	name	e:		High-adventure base participants:
Date	of b	irth:		Expedition/crew No.: or staff position:
Age:		Gender:	Height (inches):	_Weight (lbs.):
Address	S:			
				code: Phone:
				Unit leader's mobile #:
				Unit No.:
Health/	Acciden	t Insurance Company:		Policy No.:
<b>•</b>	Please	attach a photocopy of both sides of the insurance card. If you do	o not have medical insurar	ice, enter "none" above.
		ergency, notify the person below:		
Name:				Relationship:
Address:			Home phone:_	Other phone:
Alternat	e conta	ct name:		Alternate's phone:
Healt	th Hi	story		
		have or have you ever been treated for any of the following?		
Yes	No	Condition		Explain
		Diabetes	Last HbA1c percentage	ind date: Insulin pump: Yes No N
		Hypertension (high blood pressure)		
		Adult or congenital heart disease/heart attack/chest pain (angina)/ heart murmur/coronary artery disease. Any heart surgery or procedure. Explain all "yes" answers.		
		Family history of heart disease or any sudden heart-related death of a family member before age 50.		
		Stroke/TIA		
		Asthma/reactive airway disease	Last attack date:	
		Lung/respiratory disease		
		COPD		
		Ear/eyes/nose/sinus problems		
		Muscular/skeletal condition/muscle or bone issues		
		Head injury/concussion/TBI		
		Altitude sickness		
		Psychiatric/psychological or emotional difficulties		
		Neurological/behavioral disorders		
		Blood disorders/sickle cell disease		
		Fainting spells and dizziness		
		Kidney disease		
		Seizures or epilepsy	Last seizure date:	
		Abdominal/stomach/digestive problems		
		Thyroid disease		
		Skin issues		
		Obstructive sleep apnea/sleep disorders	CPAP: Yes No	
		List all surgeries and hospitalizations	Last surgery date:	



Full	name	:							ase participan		
Date	of bi	rth:									
Alle	rgies	s/Medicat									
		NEPINEPHRINE R? Exp. date (i	f ves)	☐ YES	□ NO			EAN ASTHM. Exp. date (i			YES 🗆 NO
AUTOII	102010	it: Exp. date (	1 yes/			1141	IALLIN:	Exp. dato (i	1 y 00)		_
Are you	allergic to	or do you have an	y adverse reacti	on to any of the follo	wing?						
Yes	No	Allergies or R	eactions	E	xplain	Yes	No	Allergies or	Reactions	E	xplain
		Medication Food						Plants Insect bites/stin	ige		
Liet ell	no o di o		ruped includ	ling only over the	a acustor madia	nationa		Insect bites/stil	lys .		
		-		ling any over-the			andad .	nlagge ligt or	o o oonarata ah	ant and attach	
	eckner	e if no medicat	ions are rout	пету такеп.		onal space is ne	eeaea,	piease list of	ra separate sri	eet and attach	
		Medication		Dose	Frequency				Reaso	n	
YES		O Non-pre	scription medica	tion administration is	s authorized with the	ese exceptions:					
Administ	ration of	the above medicati	ons is approved	for youth by:							
			_/ Parent/guardian s	ignature			MD/	/DO, NP, or PA sign	ature (if your state requ	ires signature)	
1						. Make sure that the	yareNO <sup>-</sup>	Texpired, include	dinginhalers and E	piPens.YouSHOL	JLDNOTSTOPtaking
V	anym	laintenance medic	cation unless ins	structed to do so by	your doctor.						
lmm	uniz	ation									
The follo	wing imn	nunizations are rec		anus immunization is					Please list any a	ndditional inform	nation about your
Yes	No No	Had Disease	ine disease colu	mn and list the date.  Immunization		Date	•		medical history		indicina do de your
			Tetanus				-(-/				
			Pertussis								
			Diphtheria								
			Measles/mu	mps/rubella							
			Polio						DO NOT WRITE		
Ħ			Chicken Pox						Review for camp or sp		
			Hepatitis A						Reviewed by:		
			Hepatitis B						Date:		
			Meningitis						Furtherapprovalrequi		No
$\overline{\Box}$			Influenza								_
			Other (i.e., HIB	)					Approved by:		
			Exemption to i	mmunizations (form	required)				Date:		



### Part C: Pre-Participation Physical

This part must be completed by certified and licensed physicians (MD, DO), nurse practitioners, or physician assistants.

Full name: _						h-adventure base participants	
Date of hirth:						edition/crew No.:	
Date of birtin.	ate of birth: or staff position:						
including	one of the nati	onal high-adve	individual has no contraindication for p enture bases, please refer to the supple ehmr to view this information online.				
Please fill in the fo	llowing info	rmation:					
Yes No						Explain	
Medical restrictions	to participate						
Yes No	Allergies or R	leactions	Explain	Yes	No	Allergies or Reactions	Explain
Foo	edication od					Plants Insectbites/stings	
Height (in	ches)		Weight(lbs.)	ВМІ		Blood Pressure	Pulse
						1	
	Normal	Abnormal	Explain Abnormalities	Examin	er's C	Certification	
Eyes						ewed the health history and examined th ng experience. This participant (with no	his person and find no contraindications for ted restrictions):
Ears/nose/throat				True	False	E	xplain
						Meets height/weight requirements.	
Lungs						Has no uncontrolled heart disease, lung	g disease, or hypertension.
Heart						Has not had an orthopedic injury, musc surgery in the last six months or posse orthopedic surgeon or treating physicia	sses a letter of clearance from his or her
Abdomen						Has no uncontrolled psychiatric disorde	ers.
						Has had no seizures in the last year.	
Genitalia/hernia						Does not have poorly controlled diabete	98.
Musculoskeletal						If planning to scuba dive, does not have	e diabetes, asthma, or seizures.
Neurological				Examiner's s			Date:
Skin issues				Address:	printou		
Other				City:		State:	ZIP code:
Guiei				Office phon	e:		

Height/Weight Restrictions
If you exceed the maximum weight for height as explained in the following chart and your planned high-adventure activity will take you more than 30 minutes away from an emergency vehicle/accessible roadway, you may not be allowed to participate.

### Maximum weight for height:

Height (inches)	Max. Weight						
60	166	65	195	70	226	75	260
61	172	66	201	71	233	76	267
62	178	67	207	72	239	77	274
63	183	68	214	73	246	78	281
64	189	69	220	74	252	79 and over	295



# High-Adventure Risk Advisory to Health-Care Providers and Parents

Phone: 305-664-4173

**Sea Base Experience.** Sea Base adventures are not risk-free. Participants must follows afety measures and take personal responsibility for their health and safety. Climate conditions include temperatures ranging from 45 to 95 degrees, high humidity, heat indexes reaching 110 degrees, and frequent, sometimes severe tropical weather. Prolonged, rigorous activities include snorkeling, scuba diving, kayaking, canoeing, sailing, hiking, and others.

It is the responsibility of participants, participant parents or guardians, participant health-care teams, and unit leaders to see that each individual—youth or adult—can safely take part in Sea Base adventures.

**Adult Participants.** It is the role of accompanying adults to ensure youth safety. Because of this, adult participants must arrive in good physical condition and have no medical conditions that could result in diverting the Sea Base staff's attention away from youth participants.

### Sea Base participants must be able to

- · Swim in a strong manner
- Climb a 6-foot ladder, unassisted, in inclement weather, from the water onto a rocking vessel
- · Self-rescue if found overboard in inclement weather

**Location.** Sea Base adventures are conducted at sea, often far from land, with limited access to emergency services. **Response times can be affected by weather, seas, and location, and can be delayed for hours.** Individuals with medical conditions that require immediate or nearly immediate access to professional medical care should not attend Sea Base.

**Right to Refuse.** Sea Base reserves the right to deny participation based on health and safety concerns and/or medical history.

### Special Needs or Medical Concerns. Any individual

with special needs or medical concerns must have an onsite advocate who understands the individual's condition and treatment and who is prepared to provide support to the individual.

**Trained Leadership.** Each crew is required to have at least one adult who is trained in wilderness first aid and CPR or has a greater professional medical certification. This leader acts as the primary first response until emergency services arrive. There are no on-site facilities for treatment or extended care at Sea Base. Sea Base does not staff professional medical personnel.

**Medications.** Individuals requiring medication should continue medications as prescribed and bring an appropriate supply. Each crew must develop a plan to secure, lock, and dispense medication.

**Allergies.** Participants with allergies that may result in severe reactions or anaphylaxis should bring an adequate supply of epinephrine auto-injectors (EpiPen) to last up to three hours.

### Florida Sea Base

Website: www.bsaseabase.org

# Recommendations Regarding Chronic Illness and/or Compromised Immune System. Persons with chronic conditions and/or

compromised immune systems should seek medical advice and education regarding medical risks associated with harsh marine environments before participating. Individuals with open wounds or who are at risk for chronic illnesses or immune disorders should not attend Sea Base.

### Hypertension (High Blood Pressure). Participants

should have a blood pressure less than 140/90. Individuals with hypertension should have the condition treated and well-controlled before attending.

### Insulin-Dependent Diabetes Mellitus. Diabetes

must be well-controlled. Hypoglycemia can lead to unconsciousness and drowning.

- Insulin-dependent persons who have been newly diagnosed or who have undergone changes in delivery systems in the last six months are advised not to participate.
- Persons with diabetes who have had frequent problems and/or hospitalizations should not participate.
- Persons using insulin to control diabetes will not be permitted to scuba dive.
- Any HbA1c test greater than 7 in the previous 12 months disqualifies a person from scuba diving.
- Persons under the age of 18 who control their diabetes with exercise and diet (without the aid of medication) and can provide three sequential hemoglobin tests with HbA1c values less than 6 may be approved to sculad dive.
- Persons over the age of 18 who control their diabetes with exercise and diet (without the aid of medication) and can provide four HbA1c tests, each with a value less than 7, within the previous 12 months may be approved to scuba dive.

**Seizures (Epilepsy).** Seizures while snorkeling or scuba diving are extremely dangerous and often fatal.

- History of loss of consciousness often precludes snorkeling or scuba diving. Formal consultation with a neurologist and/or cardiologist is required.
- No participant with a history of seizures or taking anti-epileptic medication may snorkel or scuba dive.
  - —Prospective participants with a history of infant febrile seizures may be considered for snorkeling or diving after formal consultation with a neurologist.

**Asthma.** Asthma must be well-controlled. Persons requiring use of medication and/or an inhaler must bring an ample supply.

- Persons being treated for asthma (including reactive airway disease) are disqualified from scuba diving.
- Persons with a history of asthma who have been asymptomatic and have not used medications to control asthma for five years or more may be allowed to scuba dive if resolution of asthma is specifically confirmed by their physician and includes provocative pulmonary function testing conducted by a pulmonologist.
  - —Provocative testing can include exercise, hypertonic saline, a hyperpnea test, etc.



# High-Adventure Risk Advisory to Health-Care Providers and Parents

### Florida Sea Base

Phone: 305-664-4173 Website: www.bsaseabase.org

# Recent Musculoskeletal Injuries and

**Orthopedic Surgery.** Persons with musculoskeletal problems or orthopedic surgeries within the last six months must provide a letter from their treating physician to participate.

### Psychological and Emotional Difficulties.

Any condition should be well-controlled without the services of a mental health practitioner. Participants requiring medication must bring an ample supply and take as prescribed for the duration of their trip.

- · Many psychotropic medications are not compatible with scuba diving.
- Persons taking more than one psychotropic medication will not be cleared to scuba dive.
- · Persons with anxiety will not be cleared to scuba dive.

**Weight Limits.** Participants must meet BSA height and weight guidelines. Exceptions **may** be made for individuals who do not exceed the BSA height and weight guidelines by more than 20 pounds. Due to rescue equipment weight restrictions, individuals who are 78 inches (6.5 feet) and taller cannot be offered an exception.

Scuba Participants. Persons with conditions listed as severe by the Recreational Scuba Training Council (RSTC) will not be permitted to scuba dive. Persons with conditions prohibited by BSA scuba policy will not be permitted to scuba dive. Various risk factors may exclude a person from scuba diving, either temporarily or permanently. Risk factors include, but are not limited to, ear and sinus problems, recent surgery, spontaneous pneumothorax, asthma or reactive airway disease, seizure disorders, diabetes, leukemia, sickle-cell disease, pregnancy, panic disorders, active psychosis, certain medications, and narcolepsy.

**Closing.** Sea Base, BSA is an industry leader in maritime adventures with an excellent safety record. If you have questions regarding medical policies, medical concerns, or medical approval, please contact Sea Base at 305-664-4173.













### **Diver Medical** | Participant Questionnaire

Recreational scuba diving and freediving requires good physical and mental health. There are a few medical conditions which can be hazardous while diving, listed below. Those who have, or are predisposed to, any of these conditions, should be evaluated by a physician. This Diver Medical Participant Questionnaire provides a basis to determine if you should seek out that evaluation. If you have any concerns about your diving fitness not represented on this form, consult with your physician before diving. If you are feeling ill, avoid diving. If you think you may have a contagious disease, protect yourself and others by not participating in dive training and/or dive activities. References to "diving" on this form encompass both recreational scuba diving and freediving. This form is principally designed as an initial medical screen for new divers, but is also appropriate for divers taking continuing education. For your safety, and that of others who may dive with you, answer all questions honestly.

### **Directions**

Complete this questionnaire as a prerequisite to a recreational scuba diving or freediving course.

Note to women: If you are pregnant, or attempting to become pregnant, do not dive.

I have had problems with my lungs, breathing, heart and/or blood affecting my normal physical or mental performance.	Yes □ Go to box <b>A</b>	No □
I am over 45 years of age.	Yes □ Go to box <b>B</b>	No 🗆
I struggle to perform moderate exercise (for example, walk 1.6 kilometer/one mile in 14 minutes or swim 200 meters/yards without resting), OR I have been unable to participate in a normal physical activity due to fitness or health reasons within the past 12 months.	Yes □*	No □
I have had problems with my eyes, ears, or nasal passages/sinuses.	Yes □ Go to box <b>C</b>	No □
I have had surgery within the last 12 months, OR I have ongoing problems related to past surgery.	Yes □*	No 🗆
I have lost consciousness, had migraine headaches, seizures, stroke, significant head injury, or suffer from persistent neurologic injury or disease.	Yes □ Go to box <b>D</b>	No 🗆
I am currently undergoing treatment (or have required treatment within the last five years) for psychological problems, personality disorder, panic attacks, or an addiction to drugs or alcohol; or, I have been diagnosed with a learning or developmental disability.	Yes □ Go to box <b>E</b>	No 🗆
I have had back problems, hernia, ulcers, or diabetes.	Yes □ Go to box <b>F</b>	No □
I have had stomach or intestine problems, including recent diarrhea.	Yes □ Go to box <b>G</b>	No 🗆
I am taking prescription medications (with the exception of birth control or or anti-malarial drugs other than mefloquine (Lariam).	Yes □*	No □
	I struggle to perform moderate exercise (for example, walk 1.6 kilometer/one mile in 14 minutes or swim 200 meters/yards without resting), OR I have been unable to participate in a normal physical activity due to fitness or health reasons within the past 12 months.  I have had problems with my eyes, ears, or nasal passages/sinuses.  I have had surgery within the last 12 months, OR I have ongoing problems related to past surgery.  I have lost consciousness, had migraine headaches, seizures, stroke, significant head injury, or suffer from persistent neurologic injury or disease.  I am currently undergoing treatment (or have required treatment within the last five years) for psychological problems, personality disorder, panic attacks, or an addiction to drugs or alcohol; or, I have been diagnosed with a learning or developmental disability.  I have had back problems, hernia, ulcers, or diabetes.  I have had stomach or intestine problems, including recent diarrhea.	I have had problems with my lungs, breathing, heart and/or blood affecting my normal physical or mental performance.  I am over 45 years of age.  I struggle to perform moderate exercise (for example, walk 1.6 kilometer/one mile in 14 minutes or swim 200 meters/yards without resting), OR I have been unable to participate in a normal physical activity due to fitness or health reasons within the past 12 months.  I have had problems with my eyes, ears, or nasal passages/sinuses.  I have had surgery within the last 12 months, OR I have ongoing problems related to past surgery.  I have lost consciousness, had migraine headaches, seizures, stroke, significant head injury, or suffer from persistent neurologic injury or disease.  I have lost consciousness, had migraine headaches, seizures, stroke, significant head injury, or suffer from persistent neurologic Go to box D  I am currently undergoing treatment (or have required treatment within the last five years) for psychological problems, personality disorder, panic attacks, or an addiction to drugs or alcohol; or, I have been diagnosed with a learning or developmental disability.  I have had back problems, hernia, ulcers, or diabetes.  I have had stomach or intestine problems, including recent diarrhea.  I have had stomach or intestine problems, including recent diarrhea.

# Participant Statement: I have answered all questions honestly, and understand that I accept responsibility for any consequences resulting from any questions I may have answered inaccurately or for my failure to disclose any existing or past health conditions. Participant Signature (or, if a minor, participant's parent/guardian signature required. Participant Name (Print) Participant Name (Print) Staff of Sea Base / Scouting America Instructor Name (Print) Please read and agree to the statement above by signing and dating it AND take all three pages of this form (Participant Questionnaire and the Physician's Evaluation Form) to your physician for a medical evaluation. Participation in a diving course requires your physician's approval.

Version date: 2022-02-01 1 of 3 © 2020

Participant Name		Birthdate		
------------------	--	-----------	--	--

(Print) Date (dd/mm/yyyy)

### **Diver Medical** | Participant Questionnaire Continued

BOX A – I HAVE/HAVE HAD:		
Chest surgery, heart surgery, heart valve surgery, an implantable medical device (eg, stent, pacemaker, neurostimulator), pneumothorax, and/or chronic lung disease.	Yes □*	No □
Asthma, wheezing, severe allergies, hay fever or congested airways within the last 12 months that limits my physical activity/exercise.	Yes □*	No □
A problem or illness involving my heart such as: angina, chest pain on exertion, heart failure, immersion pulmonary edema, heart attack or stroke, OR am taking medication for any heart condition.	Yes □*	No □
Recurrent bronchitis and currently coughing within the past 12 months, OR have been diagnosed with emphysema.	Yes □*	No □
Symptoms affecting my lungs, breathing, heart and/or blood in the last 30 days that impair my physical or mental performance.	Yes □*	No □
BOX B – I AM OVER 45 YEARS OF AGE AND:		
I currently smoke or inhale nicotine by other means.	Yes □*	No □
I have a high cholesterol level.	Yes □*	No □
I have high blood pressure.	Yes□*	No □
I have had a close blood relative die suddenly or of cardiac disease or stroke before the age of 50, OR have a family history of heart disease before age 50 (including abnormal heart rhythms, coronary artery disease or cardiomyopathy).	Yes □*	No □
BOX C – I HAVE/HAVE HAD:		
Sinus surgery within the last 6 months.	Yes□*	No □
Ear disease or ear surgery, hearing loss, or problems with balance.	Yes □*	No □
Recurrent sinusitis within the past 12 months.	Yes □*	No □
Eye surgery within the past 3 months.	Yes □*	No □
BOX D – I HAVE/HAVE HAD:		
Head injury with loss of consciousness within the past 5 years.	Yes □*	No □
Persistent neurologic injury or disease.	Yes □*	No □
Recurring migraine headaches within the past 12 months, or take medications to prevent them.	Yes □*	No □
Blackouts or fainting (full/partial loss of consciousness) within the last 5 years.	Yes □*	No □
Epilepsy, seizures, or convulsions, OR take medications to prevent them.	Yes □*	No □
BOX E – I HAVE/HAVE HAD:		
Behavioral health, mental or psychological problems requiring medical/psychiatric treatment.	Yes □*	No □
Major depression, suicidal ideation, panic attacks, uncontrolled bipolar disorder requiring medication/psychiatric treatment.	Yes□*	No □
Been diagnosed with a mental health condition or a learning/developmental disorder that requires ongoing care or special accommodation.	Yes □*	No □
An addiction to drugs or alcohol requiring treatment within the last 5 years.	Yes □*	No □
BOX F – I HAVE/HAVE HAD:		
Recurrent back problems in the last 6 months that limit my everyday activity.	Yes □*	No □
Back or spinal surgery within the last 12 months.	Yes □*	No □
Diabetes, either drug or diet controlled, OR gestational diabetes within the last 12 months.	Yes □*	No □
An uncorrected hernia that limits my physical abilities.	Yes □*	No □
Active or untreated ulcers, problem wounds, or ulcer surgery within the last 6 months.	Yes □*	No □
BOX G – I HAVE HAD:		
Ostomy surgery and do not have medical clearance to swim or engage in physical activity.	Yes □*	No □
Dehydration requiring medical intervention within the last 7 days.	Yes □*	No □
Active or untreated stomach or intestinal ulcers or ulcer surgery within the last 6 months.	Yes □*	No □
Frequent heartburn, regurgitation, or gastroesophageal reflux disease (GERD).	Yes □*	No □
Active or uncontrolled ulcerative colitis or Crohn's disease.	Yes□*	No □

# **Diver Medical** | Medical Examiner's Evaluation Form **Participant Name Birthdate** (Print) Date (dd/mm/yyyy) The above-named person requests your opinion of his/her medical suitability to participate in recreational scuba diving or freediving training or activity. Please visit uhms.org for medical guidance on medical conditions as they relate to diving. Review the areas relevant to your patient as part of your evaluation. **Evaluation Result** Approved - I find no conditions that I consider incompatible with recreational scuba diving or freediving. Not approved – I find conditions that I consider incompatible with recreational scuba diving or freediving. Signature of certified medical doctor or other legally certified medical provider Date (dd/mm/yyyy) **Medical Examiner's Name** (Print) **Clinical Degrees/Credentials** Clinic/Hospital **Address Phone Email** Physician/Clinic Stamp (optional) Created by the **Diver Medical Screen Committee** in association with the following bodies: The Undersea & Hyperbaric Medical Society DAN (US) **DAN Europe**

© DMSC 2020 3 of 3 10346 EN

Hyperbaric Medicine Division, University of California, San Diego



## **Diving Medical Guidance to the Physician**

These guidelines are typically used by physicians who have been approached by an individual wishing to take part in recreational scuba diving or freediving. They will usually have completed a WRSTC Diver Medical Participant Questionnaire.

Recreational scuba diving and freediving (hereafter "diving") is performed safely by many people. The risks associated with diving may be increased by certain physical conditions, and the relationship to diving may not be readily appreciated by candidates. Thus, it is important to screen divers for such conditions.

A physical examination for diving focuses on conditions that may put a diver at increased risk for decompression sickness, pulmonary overinflation with subsequent arterial gas embolization, and other conditions such as loss of consciousness, which could lead to drowning. Additionally, divers must be able to withstand some degree of thermal stress, the physiological effects of immersion, and have sufficient physical and mental reserves to deal with normal diving and possible emergencies.

The history, review of systems, and physical examination should include as a minimum the points listed below. The list of conditions that might adversely affect the diver is not exhaustive, but contains the most commonly encountered medical problems. The brief introductions serve as an alert to the nature of the risk posed.

The potential diver and his or her physician must weigh the benefits to be had by diving against an increased risk of injury or death due to the individual's medical condition. As with any recreational activity, there are limited data for diving with which to calculate the mathematical probability of injury. Experience and physiological principles only permit a qualitative assessment of relative risk.

For the purposes of this document, **Severe Risk** implies that an individual is believed to be at substantially elevated risk of injury compared with the general population. The consultants involved in drafting this document would generally discourage a candidate with such medical problems from diving. **Relative Risk** refers to a moderate increase in risk, which in some instances may be acceptable. To make a decision as to whether diving is contraindicated for this category of medical problems, physicians must base their judgment on an assessment of the individual candidate. **Temporary** Risk refers to medical problems which may preclude diving but are temporary in nature, allowing the individual to dive after they have resolved.

Following many of the sections is a short list of references that give more information on the topic. The lists are not exhaustive, but examples that may be of particular relevance.

Diagnostic studies and specialty consultations should be obtained as indicated to determine the candidate's status. A list of references is included to aid in clarifying issues that arise.

### The following sections are included in this document (click to jump to section):

**BEHAVIORAL HEALTH CARDIOVASCULAR SYSTEMS GASTROINTESTINAL HEMATOLOGICAL** METABOLIC AND ENDOCRINOLOGICAL **NEUROLOGICAL ORTHOPEDIC** 

**OTOLARYNGOLOGICAL** 

**PULMONARY** 

### **BEHAVIORAL HEALTH**

Behavioral health is one of the most difficult aspects of diver candidate evaluation, because many relevant potential problems may not be apparent and are not easily assessed in an office consultation. This is also an aspect of evaluating suitability for diving in which the diving instructor, who observes the candidate in the field, must also play a part.

The diving candidate must be capable of learning and applying a theoretical knowledge base for diving. Significant intellectual handicap is incompatible with independent diving.

Motivational and behavioral traits should be considered if there is obvious related history or problems become apparent during training. Candidates who appear unmotivated, irresponsible, or prone to distraction or panic should be discouraged from diving.

A history of psychiatric disease is not in and of itself disqualifying. Psychotropic medications can be problematic if they are associated with altered level of awareness or sedation, or may alter seizure threshold, (e.g., benzodiazepines, narcotics). What is of primary importance is the individual's current psychological state, and anticipated impact of their mental/psychological history relative to their ability to navigate the potential and anticipated challenges and stresses of diving. The level of baseline mental health, with or without medication, is therefore of greater importance than the theoretical effects of a given medication or class of medications while diving.

Candidates with major depression, bipolar disorder, psychoses, or current drug or alcohol abuse should not dive. Even if a candidate is well controlled on medication (see below for discussion of SSRIs), there may be risks associated with the use of potent antidepressant and antipsychotic drugs in the underwater environment. The tendency for potent psychotropic drugs to impair concentration and cause drowsiness is of particular concern, as is their potential to lower the seizure threshold, and the lack of research data evaluating potential interactions with the pressure environment. Candidates with a past history of major psychiatric problems or drug/alcohol abuse who are stable without medication and withdrawn from drugs and alcohol can be considered on a case-by-case basis, preferably by a physician trained in diving medicine.

Perhaps the most challenging group of candidates from a behavioral perspective in the modern context is those with "mild" depression (those who have never been hospitalized for psychiatric treatment or placed on psychiatric hold or attempted self-harm) or those with mood disturbances treated with selective serotonin reuptake inhibitors (SSRIs). The general use of SSRIs has increased dramatically over recent years in many countries. There are no data describing use of SSRIs among divers, but anecdotally the numbers are significant. Concerns over diving while using SSRIs relate to the disorder being treated and to the potential interaction between the drug and diving. There are many candidates taking these drugs whose mild mood disturbance would not of itself constitute a reason to avoid diving. Evaluation of the potential for an interaction between SSRIs and diving is more difficult. There are no published reports of apparent problems despite what is almost certainly a large number of divers using them. Diving while taking an SSRI is probably acceptable provided that: the treated mood disturbance was mild prior to treatment and has been well controlled by the drug; the drug has been used for at least one month without evidence of relevant side effects; and the candidate is fully counseled about (and accepting of) the relevant risks. If the candidate is considering diving beyond the traditional recreational envelope or using gases other than air, he or she should consult an appropriate diving medicine specialist.

There are also potential risks associated with other drugs used to treat psychiatric conditions, including serotoninnorepinephrine reuptake inhibitors (SNRIs), tricyclic antidepressants (TCAs), monoamine oxidase inhibitors (MAOIs), and atypical agents (including bupropion). Candidates on these medications should be evaluated on a case-by-case basis.

### **Severe Risk Conditions**

- Active major depression, bipolar or psychotic disorder
- History of panic attacks
- Drug or alcohol abuse
- Severe intellectual handicap

### **Relative Risk Conditions**

- Questionable motivation to dive solely to please spouse, partner or family member, or to prove oneself in the face of personal fears
- Developmental delay/Cognitive impairment
- Anxiety disorder
- History of drug or alcohol abuse
- History of major depression, bipolar, or psychotic disorder
- Use of psychotropic medications
- Claustrophobia or agoraphobia

### CARDIOVASCULAR SYSTEMS

Diving places increased demands on the heart. Immersion itself results in an increase in cardiac preload, as does peripheral vasoconstriction with an increase in blood pressure. These changes are typically accompanied by sustained mild to moderate exercise. Perhaps not surprisingly, almost 30% of recreational diving fatalities have a cardiac event as the disabling injury. It follows that the primary goals of evaluating the cardiovascular system in a diving candidate are to identify those who appear to be at risk of myocardial ischemic events, myocardial insufficiency, or other cardiac events (such as arrhythmias) that might disable a diver underwater, and to establish that the candidate has an adequate exercise capacity for diving.

With the above in mind, some cardiac diagnoses are considered to render a candidate unsuitable for diving, including: untreated symptomatic coronary artery disease, dilated or obstructive or previous stress cardiomyopathy, congestive heart failure, moderate or worse pulmonary hypertension, long QT syndrome or other arrhythmia-inducing channelopathies, paroxysmal arrhythmias causing unconsciousness or impairment of exercise capacity, poor exercise capacity of apparent cardiac origin, moderate to severe valvular lesions, complex congenital cardiac disease, atrial septal defect, and the presence of an implanted cardiac defibrillator.

Potential candidates with any of the following should be investigated to exclude a disqualifying condition:

- Exertional chest pain, dyspnea, palpitations, or syncope
- Unexplained syncope/near syncope
- Heart murmur
- Hypertension
- Family history of premature death (sudden/unexpected or cardiac) before age 50, cardiac disease before age 50, cardiomyopathy, arrhythmia, or channelopathy

It is strongly recommended that these candidates be evaluated in consultation with a physician trained in diving medicine and possibly a cardiologist. Successful treatment of disqualifying cardiac disorders may result in a candidate becoming suitable for diving. For example, a candidate with coronary artery disease (including previous myocardial infarction) who has been successfully revascularized may be suitable for diving if inducible ischemia can be excluded and adequate exercise capacity demonstrated (for example, in an exercise stress test). The capacity to sustain exercise at 6 MET (metabolic equivalent of task; 1 MET approximates resting metabolic rate, assumed to approximate an oxygen consumption of 3.5 mL/kg/min; 6 MET approximates an effort of six times resting metabolic rate, approximating an oxygen consumption of 21 mL/kg/min is a pragmatic expectation for a recreational diver, but there may be an occasional need to exercise transiently at higher levels during diving. Similarly, a candidate with a history of paroxysmal arrhythmia who has undergone successful pathway ablation may be suitable for diving. Candidates with any of the above diagnoses who wish to consider diving after appropriate treatment are best referred to a physician trained in diving medicine for evaluation.

Asymptomatic candidates over 45 years of age with risk factors for coronary artery disease should undergo evaluation by a physician. Individuals with a predicted 5-10 year risk of a cardiovascular event >10% using a cardiac risk calculator should be investigated for coronary disease unless they provide a credible history of exercise capacity which renders significant coronary disease very unlikely. A coronary calcium score is a suitable initial investigation, and a myocardial perfusion scan, stress echocardiogram, or CT coronary angiogram should be considered in following up a positive calcium score. Consideration of a tailored investigation pathway for the individual diving candidate is ideally undertaken by a cardiologist in consultation with a physician trained in diving medicine. Candidates who prove to have inducible ischemia or obstructive lesions justifying intervention should not dive until completion of the intervention and demonstration of its success. Candidates with non-obstructive coronary disease not requiring invasive intervention should have aggressive management of risk factors and may be suitable for diving if adequate exercise capacity can be demonstrated. Although an exercise ECG is relatively insensitive to early coronary disease, it has the advantage of demonstrating exercise capacity and can be modified to test sustained exercise at 6 MET.

Left ventricular hypertrophy (LVH) is a risk factor for arrhythmias, which may be induced by exercise or immersion. Candidates for diving with this condition should be counseled about the risks of diving.

A patent foramen ovale (PFO) that exhibits right-to-left shunting with no or minimal provocation is a risk factor for serious neurological decompression sickness. In established divers, such lesions are usually discovered by bubble contrast echocardiography conducted after a relevant episode of decompression sickness. These divers are usually advised either to cease diving, modify their diving to reduce venous bubble formation (venous bubbles crossing from right to left are almost certainly the vectors of harm in this setting), or to have the PFO repaired. Occasionally, new diver candidates have a previously discovered PFO, and in such cases an objective assessment of the shunting behavior of the lesion is required in order to adequately counsel the candidate about the implied risks in diving. If not already done, this is best achieved using bubble contrast transthoracic echocardiography at rest and with provocative maneuvers. It is strongly recommended that the results of such tests are discussed with a physician trained in diving medicine. Routine screening of all diving candidates for PFO is not recommended.

In relation to some specific cardiovascular diagnoses: Treated hypertension with adequate control is acceptable in diving in the absence of other risk factors that would meet a risk threshold indicating screening for coronary artery disease. Atrial fibrillation that is adequately rate-controlled in a candidate without inducible myocardial ischemia and who exhibits adequate exercise capacity is acceptable in diving. However, many such candidates are anticoagulated and the risks of diving whilst anticoagulated would need to be understood and carefully considered by the candidate. This is best achieved through discussion with a physician trained in diving medicine.

Immersion pulmonary edema is a problem that has been seen in swimmers, compressed gas divers, and freedivers. The condition may be under-diagnosed. Risk factors include hypertension, valvular disease, diastolic dysfunction, cardiomyopathies, pulmonary hypertension, hyperhydration, immersion, cold stress, constrictive garments, exercise, and for compressed gas divers, increased breathing resistance (affected by equipment, gas density, and body position), and for freedivers, pulmonary squeeze due to compression during descent. A single episode of immersion pulmonary edema may contra-indicate further diving if no modifiable risk factors are found. Repetitive cases represent a strong contra-indication. A diver or new diving candidate with such a history should be referred to a physician trained in diving medicine for discussion of the relevant issues.

Candidates with pacemakers may be able to dive, though pacemaker-dependent candidates should consider the risks carefully. The pathologic process that necessitated the pacemaker should be considered as should the candidate's functional capacity (see above). Pacemakers must be certified by the manufacturer as able to withstand the pressure changes involved in recreational diving. Devices vary in this regard, but diving beyond 30 meters/100 feet with any of them is unwise.

### **Severe Risk Conditions**

- Untreated symptomatic coronary artery disease
- Dilated or obstructive cardiomyopathy
- Heart failure
- Pulmonary hypertension
- Long QT syndrome or other arrhythmia-inducing channelopathies
- Paroxysmal arrhythmias causing unconsciousness or impairment of exercise capacity
- Poor exercise capacity of apparent cardiac origin
- Moderate to severe valvular lesions
- Complex congenital cardiac disease
- Atrial septal defect
- Presence of an implanted cardiac defibrillator
- Multiple episodes of immersion pulmonary edema

### **Relative Risk Conditions**

- Treated coronary artery disease
- Collectively, risk factors such as age >45 years, hypertension, smoking, elevated cholesterol and a positive family history may indicate investigation for coronary artery disease
- History of dysrhythmias requiring medication for suppression
- Mild valvular lesions (need periodic re-evaluation)
- Cardiac prostheses or arrhythmias requiring anticoagulation
- Pacemakers
- Single previous episode of immersion pulmonary edema
- Marfan syndrome or other connective tissue disorder (severe risk if there is a history of dissection)
- Left ventricular hypertrophy

### References

Denoble PJ, Holm JR, eds. Patent Foramen Ovale and Fitness to Dive Consensus Workshop Proceedings. Durham, NC: Divers Alert Network, 2015; 160 pp.

Kumar M, Thompson PD. A literature review of immersion pulmonary edema. Physic Sportsmed. 2018; 47(2):148-151.

Lafay V, Trigano JA, Gardette B, Micoli C, Carre F. Effects of hyperbaric exposures on cardiac pacemakers. *Br J Sports Med.* 2008;42(3):212-216

Mitchell SJ, Bove AA. Medical screening of recreational divers for cardiovascular disease: Consensus discussion at the Divers Alert Network Fatality Workshop. Undersea Hyperb Med. 2011; 38(4), 289-296.

Moon RE, Bove AA, Mitchell SJ. PFO statement. In: Denoble PJ, Holm JR. eds. Patent Foramen Ovale and Fitness to Dive Consensus Workshop Proceedings. Durham, NC: Divers Alert Network, 2016; 156-160.

Pollock NW. Aerobic fitness and underwater diving. Diving Hyperb Med. 2007; 37(3): 118-124.

Smart D, Mitchell SJ, Wilmshurst P, Turner M, Banham N. Joint position statement on persistent (patent) foramen ovale and diving. South Pacific Underwater Medicine Society (SPUMS) and the United Kingdom Sports Diving Medical Committee (UKSDMC). Diving Hyperb Med. 2015; 45(2), 129-131.

### GASTROINTESTINAL

In general terms, there should be no gastrointestinal conditions present that increase the likelihood of vomiting, reflux, bleeding, perforation, diarrhea, or pain. Altered anatomical relationships secondary to surgery or malformations that lead to gas trapping may cause serious problems. Trapped gas expands as the diver surfaces and can lead to rupture or, in the case of the upper GI tract, emesis. Emesis underwater may lead to drowning. Dive activities may take place in areas remote from medical care, and the possibility of acute recurrences of disease must be considered.

### **Severe Risk Conditions**

- Active inflammatory bowel disease
- Gastric outlet obstruction of a degree sufficient to produce recurrent vomiting
- Chronic or recurrent small bowel obstruction
- Severe gastroesophageal reflux
- Achalasia
- Paraesophageal hernia
- Gastroparesis

### **Relative Risk Conditions**

- Inflammatory bowel disease when guiescent
- Functional bowel disorders

### **Temporary Risk Conditions**

- Peptic ulcer disease associated with pyloric obstruction or severe reflux
- Unrepaired hernias of the abdominal wall large enough to contain bowel within the hernia sac could incarcerate

### References

Bennett PB, Cronje FJ, Campbell E, Marroni A, Pollock NW. Assessment of Diving Medical Fitness for Scuba Divers and Instructors. Flagstaff, AZ: Best Publishing. 2006; 241 pp.

Vote D. Gastrointestinal issues – consider them before returning to diving. <a href="https://www.diversalertnetwork.org/medical/articles/">https://www.diversalertnetwork.org/medical/articles/</a> Gastrointestinal Issues

US Navy Diving Manual, Volume 2, Revision 7. Gastrointestinal distension.NAVSEA 0910-LP-115-1921. Naval Sea Systems Command: Washington, DC, 2016: 3-31-3-32.

### **HEMATOLOGICAL**

Abnormalities resulting in altered rheological properties may theoretically increase the risk of decompression sickness. Bleeding disorders could worsen the effects of otic or sinus barotrauma and exacerbate the injury associated with inner ear or spinal cord decompression sickness. Spontaneous bleeding into the joints (eg, in hemophilia) may be difficult to distinguish from decompression illness. Thrombophilic disorders (hereditary or acquired) may facilitate vascular thrombosis and susceptibility to DCS.

### **Relative Risk Conditions**

- Sickle cell disease
- Polycythemia vera
- Leukemia
- Hemophilia/Impaired coagulation
- Recent blood transfusion
- Recent thrombotic episodes
- Hereditary hypercoagulability conditions
  - Factor V Leiden
  - Prothrombin 20210A
  - Protein C deficiency
  - Protein S deficiency
  - Antithrombin deficiency

### **Temporary Risk Conditions**

Prescription of anti-coagulant drugs of any kind, including platelet aggregation inhibitors

### References

Bennett PB, Cronje FJ, Campbell E, Marroni A, Pollock NW. Assessment of Diving Medical Fitness for Scuba Divers and Instructors. Flagstaff, AZ: Best Publishing. 2006; pp 97-104.

Parker J. Haematology. In: The Sports Diving Medical, 2nd Edition. JL Publications, Melbourne 2002, pp 100-102.

Wendling J, et al. Haematological disorders. In: Medical Assessment of Fitness to Dive. International Edition. Hyperbaric Editions CH 2502 Biel, 2001, pp 126. ISBN 3-9522284-1-9.

### METABOLIC AND ENDOCRINOLOGICAL

States of altered hormonal or metabolic function should be assessed according to their impact on the individual's ability to tolerate the moderate exercise requirement and environmental stress of sport diving. Obesity may predispose the individual to decompression sickness, can impair exercise tolerance and is a risk factor for coronary artery disease.

### **Severe Risk Conditions**

- The potentially rapid change in level of consciousness associated with hypoglycemia in diabetics on insulin therapy or certain oral hypoglycemic medications can result in drowning. Diving is therefore generally contraindicated, except when conducted according to the consensus guidelines for recreational diving with diabetes.
- Pregnancy: The effect of venous emboli formed during decompression on the fetus has been proven to be potentially detrimental to fetus health. Diving is therefore not recommended during any stage of pregnancy or for women actively seeking to become pregnant. (Note that in cases where pregnancy is discovered after diving, it is not considered grounds for termination.)

### **Relative Risk Conditions**

- Hormonal excess or deficiency
- Obesity
- Renal insufficiency

### References

Damnon F, de Rham M, Baud D. Should a pregnancy test be required before scuba diving? Br J Sports Med. 2016; 50(18): 1159-1160.

Dear GdeL, Pollock NW, Uguccioni DM, Dovenbarger J, Feinglos MN, Moon RE. Plasma glucose response to recreational diving in divers with insulin-requiring diabetes. Undersea Hyperb Med. 2004; 31(3): 291-301.

Held HE, Pollock NW. The risks of diving while pregnant - reviewing the research. Alert Diver. 2007; Mar/Apr: 48-51.

Pollock NW, Uguccioni DM, Dear GdeL. Diabetes and recreational diving: guidelines for the future. Diving Hyperb Med 2006; 36(1): 29-34.

### **NEUROLOGICAL**

Neurological illnesses, especially those affecting the spinal cord and peripheral nerves, should be assessed according to the degree of functional compromise present. Any condition that diminishes the reserve capacity of the spinal cord may reduce the likelihood of a full functional recovery, should an episode of spinal decompression sickness occur. Conditions in which there can be a waxing and waning of neurological symptoms and signs, such as migraine or demyelinating disease, may contraindicate diving, because an exacerbation or attack of the pre-existing disease (eg, migraine headache with aura) may be difficult to distinguish from neurological decompression sickness. A history of head injury resulting in unconsciousness should be evaluated for risk of seizure. A diagnosis of epilepsy is considered an absolute contraindication for diving.

### **Severe Risk Conditions**

Any abnormalities where there is a significant probability of unconsciousness, hence putting the diver at increased risk of drowning. Divers with spinal cord or brain abnormalities where perfusion is impaired may be at increased risk of decompression sickness.

Some conditions are as follows:

- Epilepsy or history of seizures, other than childhood febrile seizures
- History of transient ischemic attack (TIA) or cerebrovascular accident (CVA)
- History of serious (central nervous system, cerebral or inner ear) decompression sickness with residual deficits
- Recurrent episodes of loss of consciousness or fainting

### **Relative Risk Conditions**

Complicated migraine headaches, particularly if severe, frequent or presenting with neurological manifestations eg, motor, sensory or cognitive disturbance.

- History of head injury with sequelae other than seizure
- Herniated nucleus pulposus
- Intracranial tumor or aneurysm
- Peripheral neuropathy
- Multiple sclerosis
- Trigeminal neuralgia
- History of spinal cord or brain injury
- Parkinson's disease

### References

Bennett PB, Cronje FJ, Campbell E, Marroni A, Pollock NW. Assessment of Diving Medical Fitness for Scuba Divers and Instructors. Flagstaff, AZ: Best Publishing. 2006; 241 pp. 173-188.

Burkett JG, Nahas-Geiger SJ. Diving Headache. Curr Pain Headache Rep. 2019;23(7):46.

Massey EW, Moon RE. Neurology and diving. Handb Clin Neurol. 2014;120:959-969.

Rosinska J, Łukasik M, Kozubski W. Neurological complications of underwater diving. Neurol Neurochir Pol. 2015;49(1):45-51.

UK Diving Medical Committee, Neurological disease. http://www.ukdmc.org/medical-conditions/neurological-disease/

### **ORTHOPEDIC**

Mobility above and under the water is an essential requirement for any sport or recreational diver. Entering the water from shore or a dive boat, underwater propulsion and exiting into a dive boat or onto shore should be possible without great difficulty.

Relative impairment of mobility, particularly on a boat or ashore with equipment weighing up to 30 kg/66 lb (or significantly more in the case of cold water or for more equipment intensive activities, for example), must be assessed. Orthopedic conditions of a degree sufficient to impair exercise performance may increase the risk.

In some cases, like amputations resulting in various degrees of disability, it would be advisable to judge case by case by a physician trained in diving medicine.

### **Relative Risk Conditions**

- Amputation
- Scoliosis: must also assess impact on respiratory function and exercise performance

- Aseptic necrosis: possible risk of accelerated progression due to the effects of decompression
- Disc prolapse
- Habitual luxation (eg, shoulder, hip, patella)
- Degenerative joint diseases

### **Temporary Risk Conditions**

- Back pain
- Fractures until complete healing of bone and soft tissue and positive weight bearing tests taking into consideration the weight of the used dive gear on land
- Muscle-tendon and ligament injuries
- Completion of physiotherapy/rehabilitation regimes

### References

Moeller JL. Contraindications to athletic participation. Physic Sportsmed. 1996; 24(9): 57-75.

### **OTOLARYNGOLOGICAL**

Equalization of pressure must take place during ascent and descent between ambient water pressure and the external auditory canal, middle ear, and paranasal sinuses. Failure of this to occur results at least in pain and in the worst-case rupture of the occluded space with disabling and possible lethal consequences. The inner ear is fluid filled and therefore noncompressible. The flexible interfaces between the middle and inner ear, the round and oval windows are, however, subject to pressure changes. Previously ruptured but healed round or oval window membranes may be prone to reinjury with marked overpressurization during vigorous or explosive Valsalva maneuvers. The larynx and pharynx must be free of obstruction to airflow. The laryngeal and epiglottic structures must function normally to prevent aspiration. Mandibular and maxillary function must be capable of allowing the candidate to hold a scuba regulator mouthpiece. Individuals who have had mid-face fractures may be prone to barotrauma and rupture of the air-filled cavities involved.

### **Severe Risk Conditions**

- Monomeric tympanic membrane (TM)
- Open TM perforation
- Tube myringotomy
- History of stapedectomy
- History of ossicular chain surgery
- History of inner ear surgery
- Facial nerve paralysis secondary to barotrauma
- Inner ear disease other than presbycusis
- Uncorrected upper airway obstruction
- Laryngectomy or status post partial laryngectomy
- Tracheostomy
- Uncorrected laryngocele
- History of vestibular decompression sickness
- Symptomatic nasal or sinus polyps
- Ménière's disease

### **Relative Risk Conditions**

- Recurrent otitis externa
- Significant obstruction of external auditory canal

- History of significant cold injury to pinna
- Eustachian tube dysfunction
- Recurrent otitis media or sinusitis
- History of TM perforation
- History of tympanoplasty
- History of mastoidectomy
- Significant conductive or sensorineural hearing impairment
- Facial nerve paralysis not associated with barotrauma
- Full prosthodontic devices
- History of mid-face fracture
- Unhealed oral surgery sites
- History of head and/or neck therapeutic radiation
- History of temporomandibular joint dysfunction
- History of round window rupture
- Symptomatic nasal septum deviation
- Recurrent benign positional vertigo
- Otosclerosis

### References

Lechner M, Sutton L, Fishman JM, Kaylie DM, Moon RE, Masterson L, et al. Otorhinolaryngology and diving - part 1: otorhinolaryngological hazards related to compressed gas scuba diving: a review. JAMA Otolaryngol Head Neck Surg. 2018;144(3):252-258.

Lechner M, Sutton L, Fishman JM, Kaylie DM, Moon RE, Masterson L, et al. Otorhinolaryngology and diving – part 2: otorhinolaryngological fitness for compressed gas scuba diving: a review. JAMA Otolaryngol Head Neck Surg. 2018;144(3):259-263.

Molvaer OI. Otorhinolaryngological aspects of diving. In: Bennett PB, Elliott DH, eds. Physiology and Medicine of Diving, 5th ed. Saunders, Edinburgh, 2003. P227-P264.

Wendling J, et al. Otorhinolaryngology. In: Medical Assessment of Fitness to Dive. International Edition. Hyperbaric Editions CH 2502 Biel, 2001. Pp25-48. ISBN 3-9522284-1-9.

### **PULMONARY**

Any process or lesion that impedes airflow from the lungs places the diver at risk for pulmonary over inflation with alveolar rupture and the possibility of cerebral air embolization. Many interstitial diseases predispose to spontaneous pneumothorax: asthma, chronic obstructive pulmonary disease (COPD), cystic or cavitating lung diseases may all cause air trapping.

Undersea and Hyperbaric Medical Society and British Thoracic Society guidelines recommend that asthmatics should be advised not to dive if they have wheeze precipitated by exercise, cold, or emotion. Asthmatic individuals who are currently well controlled and have normal pulmonary function tests may dive if they have a negative exercise test. Many people with asthma have well controlled disease and are physically fit. They may, however, show minor abnormalities on spirometry at rest or after exercise. Those with a history of severe or unpredictable acute exacerbations are not fit to dive. For those without such a history, the overriding consideration is that the candidate must be physically fit and not impaired after exercise or cold air breathing, which is the normal case of gas expanding from within a scuba cylinder. The best way to assess fitness is with an exercise test. Inhalation challenge tests (eg, using histamine, hypertonic saline or methacholine) are not sufficiently standardized to be interpreted in the context of scuba diving. If persons with breathing issues are cleared to dive, they need to take their regular inhalers and should not dive if suffering symptoms suggestive of exacerbation. Note that the FEV1/FVC ratio may be reduced below predicted, but provided there is no deterioration after exercise and the person performs well on the exercise test, a mildly obstructed spirometric tracing on its own is not a contraindication to diving.

A pneumothorax that occurs while diving may be catastrophic. As the diver ascends, trapped gas expands and could produce a tension pneumothorax. In addition to the risk of pulmonary barotrauma, respiratory disease due to either structural disorders of the lung or chest wall or neuromuscular disease may impair exercise performance. Individuals who have experienced spontaneous pneumothorax are at risk of recurrence, and should avoid diving, even after a surgical procedure designed to prevent recurrence (such as pleurodesis). Surgical procedures either do not correct the underlying lung abnormality (eg, pleurodesis, apical pleurectomy) or may not totally correct it (eg, resection of blebs or bullae). A high-resolution CT (HRCT) scan of the lungs may reveal cysts or blebs that represent a risk. Persons who have no parenchymal abnormality on HRCT and have had bilateral surgical pleurodesis (including VATS pleurodesis) may be cleared to dive. However, in most cases, a history of spontaneous pneumothorax will be an absolute contraindication to diving. Traumatic pneumothorax is not a problem as the likelihood of subsequent spontaneous pneumothorax is vanishingly low.

Structural disorders of the chest or abdominal wall or neuromuscular disorders may impair cough, which could be life threatening if water is aspirated. Respiratory limitation due to disease is compounded by the combined effects of immersion (causing a restrictive deficit) and the increase in gas density, which increases in proportion to the ambient pressure (causing increased airway resistance). Formal exercise testing may be helpful.

The emergence of COVID-19 has placed an additional layer of complexity related to fitness to dive evaluations. It is beyond the scope of this document to prescribe or mandate specific tests or timelines related to fitness to dive determinations. What is of importance is awareness of the potential body systems effected by COVID-19, and to take a thoughtful and thorough history related to disease course, time since the infection resolved, and state of physical and mental health at the time of the examination.

Clinical factors that are important to consider include symptom severity during the infection and need for intensive care (e.g., ventilator support). Disease severity likely correlates with the extent of pulmonary injury and potential cardiac involvement, and in the case of intubation, may be associated with severe deconditioning, muscle atrophy and even post-traumatic stress. As such, assessment of the diver with a history of COVID-19, may require more than just a pulmonary evaluation. At the time of this publication, the medical community does not have sufficient data to support arbitrary requirements for specific testing, nor duration of post-infection convalescence after which individuals can be considered safe to return to diving.

The following documents provide current guidance on investigation of COVID-19 patients prior to diving. This is an area that is evolving and updated often; please see these resources for more current information and considerations regarding these issues.

UC San Diego Guidelines for Evaluation of Divers during COVID-19 pandemic

Centers for Disease Control and Prevention, People Who Are at Higher Risk for Severe Illness

<u>European Committee for Hyperbaric Medicine and European Underwater and Baromedical Society, COVID-19 Pandemic – Position Statements</u>

For those looking for aseptic practices, the following resources may be useful:

Divers Alert Network Europe

**Divers Alert Network Americas** 

### **Severe Risk Conditions**

- History of spontaneous pneumothorax (see notes)
- Impaired exercise performance due to respiratory disease
- Respiratory impairment secondary to cold gas breathing
- Pulmonary hypertension

### **Relative Risk Conditions**

- Asthma, reactive airway disease (RAD), exercise-induced bronchospasm (EIB) or COPD (see notes)
- Solid, cystic or cavitating lesion
- Pneumothorax secondary to:
  - Thoracic surgery
  - Trauma or pleural penetration (see notes)
  - Previous overinflation injury
- Obesity
- History of immersion pulmonary edema or restrictive disease
- Interstitial lung disease: may increase the risk of pneumothorax and likely to limit exertion
- Sleep apnea

### References

Godden D, Currie G, Denison D, Farrell P, Ross J, Stephenson R, Watt S, Wilmshurst P. British Thoracic Society guidelines on respiratory aspects of fitness for diving. Thorax. 2003;58:3-13.

### **DIVERS ALERT NETWORK (DAN)**

Divers Alert Network (DAN), a non-profit organization, provides medical information and advice for the benefit of the diving public. DAN is not a regulatory agency and does not set physical standards or guidelines for scuba diving. The responsibility for the decision of whether or not to dive is generally left up to the individual, the physician, as well as the dive provider. This decision, however, should be based on the most current diving medical information available.

DAN may be able to provide current medical literature and information that can be used to assist in this decision-making process. If desired, DAN may also provide referrals to local physicians who are knowledgeable in dive medicine and physiology. However, DAN cannot and does not decide whether an individual may or may not participate in the sport of scuba diving. For more information, please feel free to contact one of the DAN offices listed below.

### DAN (US)

Physicians and other medical professionals associated with DAN are available for consultation by phone, during normal business hours Monday through Friday, 8:30 AM to 5:00 PM Eastern Time US.

+1-919-684-2948 ext. 6222

www.DAN.org

### **DAN Europe (Italy)**

+39-085-8930333

www.DANEurope.org

### **DAN World (Australia)**

+61-3-9886-9166

www.DANAP.org

### **DAN Southern Africa (South Africa)**

+27-11-266-4900

www.DANSA.org

### **DAN Japan (Yokohama)**

+045-228-3066 Medical Information Line service is provided in Japanese only. www.danjapan.gr.jp



### Non-Agency Disclosure and Acknowledgment Agreement

In European Union and European Free Trade Association countries use alternative form.

### Please read carefully and fill in all blanks before signing.

### Liability Release and Assumption of Risk Agreement

In European Union and European Free Trade Association countries use alternative form.

D.I		c 11		C:11						r		
PIESCE	read	carefully	and	TILL	ın	all	hlan	kς	hei	tore	SIC	nına
· icasc	LCUU	carcially	ullu			all	Didii	1/2	2	010	219	9

I, \_\_\_\_\_\_, hereby affirm that I am aware that skin and scuba diving have inherent risks which may result in serious injury or death.

I understand that diving with compressed air involves certain inherent risks; including but not limited to decompression sickness, embolism or other hyperbaric/air expansion injury that require treatment in a recompression chamber. I further understand that the open water diving trips which are necessary for training and for certification may be conducted at a site that is remote, either by time or distance or both, from such a recompression chamber. I still choose to proceed with such instructional dives in spite of the possible absence of a recompression chamber in proximity to the dive site.

I understand and agree that neither my instructor(s), Staff of Sea Base/ Scouting America,

the facility through which I receive my instruction,

Sea Base / Scouting America,

nor PADI Americas, Inc., nor its affiliate and subsidiary corporations, nor any of their respective employees, officers, agents, contractors or assigns (hereinafter referred to as "Released Parties") may be held liable or responsible in any way for any injury, death or other damages to me, my family, estate, heirs or assigns that may occur as a result of my participation in this diving program or as a result of the negligence of any party, including the Released Parties, whether passive or active.

In consideration of being allowed to participate in this course (and optional Adventure Dive), hereinafter referred to as "program," I hereby personally assume all risks of this program, whether foreseen or unforeseen, that may befall me while I am a participant in this program including, but not limited to, the academics, confined water and/or open water activities.

I further release, exempt and hold harmless said program and Released Parties from any claim or lawsuit by me, my family, estate, heirs or assigns, arising out of my enrollment and participation in this program including both claims arising during the program or after I receive my certification.

I also understand that skin diving and scuba diving are physically strenuous activities and that I will be exerting myself during this program, and that if I am injured as a result of heart attack, panic, hyperventilation, drowning or any other cause, that I expressly assume the risk of said injuries and that I will not hold the Released Parties responsible for the same.

I further state that I am of lawful age and legally competent to sign this liability release, or that I have acquired the written consent of my parent or guardian. I understand the terms herein are contractual and not a mere recital, and that I have signed this Agreement of my own free act and with the knowledge that I hereby agree to waive my legal rights. I further agree that if any provision of this Agreement is found to be unenforceable or invalid, that provision shall be severed from this Agreement. The remainder of this Agreement will then be construed as though the unenforceable provision had never been contained herein.

I understand and agree that I am not only giving up my right to sue the Released Parties but also any rights my heirs, assigns, or beneficiaries may have to sue the Released Parties resulting from my death. I further represent I have the authority to do so and that my heirs, assigns, or beneficiaries will be estopped from claiming otherwise because of my representations to the Released Parties.

I, Participant Name

BY THIS INSTRUMENT AGREE TO EXEMPT AND RELEASE MY INSTRUCTORS, Staff of Sea Base/ Scouting America

THE FACILITY THROUGH WHICH I RECEIVE MY INSTRUCTION, Sea Base / Scouting America , AND PADI AMERICAS, INC., AND ALL RELATED ENTITIES AS DEFINED

PADI AMERICAS, INC., AND ALL RELATED ENTITIES AS DEFINED ABOVE, FROM ALL LIABILITY OR RESPONSIBILITY WHATSOEVER FOR PERSONAL INJURY, PROPERTY DAMAGE OR WRONGFUL DEATH HOWEVER CAUSED, INCLUDING, BUT NOT LIMITED TO, THE NEGLIGENCE OF THE RELEASED PARTIES, WHETHER PASSIVE OR ACTIVE.

I HAVE FULLY INFORMED MYSELF AND MY HEIRS OF THE CONTENTS OF THIS NON-AGENCY DISCLOSURE AND ACKNOWLDGE-MENT AGREEMENT AND LIABILITY RELEASE AND ASSUMPTION OF RISK AGREEMENT BY READING BOTH BEFORE SIGNING RELOW ON REHALF OF MYSELF AND MY HEIRS.

D	ELOW ON BEHALF OF MITSELF AND MIT HEIRS.	
	Participant's Signature	Date (Day / Month / Year)
_	Signature of Parent or Guardian (where applicable)	Date (Day / Month / Year)



### Standard Safe Diving Practices Statement of Understanding

### Please read carefully before signing.

This is a statement in which you are informed of the established safe diving practices for skin and scuba diving. These practices have been compiled for your review and acknowledgement and are intended to increase your comfort and safety in diving. Your signature on this statement is required as proof that you are aware of these safe diving practices. Read and discuss the statement prior to signing it. If you are a minor, this form must also be signed by a parent or guardian.

- 1. Maintain good mental and physical fitness for diving. Avoid being under the influence of alcohol or dangerous drugs when diving. Keep proficient in diving skills, striving to increase them through continuing education and reviewing them in controlled conditions after a period of diving inactivity, and refer to my course materials to stay current and refresh myself on important information.
- Be familiar with my dive sites. If not, obtain a formal diving orientation from a knowledgeable, local source. If diving conditions are worse than those in which I am experienced, postpone diving or select an alternate site with better conditions. Engage only in diving activities consistent with my training and experience. Do not engage in cave or technical diving unless specifically trained to do so.
- 3. Use complete, well-maintained, reliable equipment with which I am familiar; and inspect it for correct fit and function prior to each dive. Have a buoyancy control device, low-pressure buoyancy control inflation system, submersible pressure gauge and alternate air source and dive planning/monitoring device (dive computer, RDP/dive tables—whichever you are trained to use) when scuba diving. Deny use of my equipment to uncertified divers.
- 4. Listen carefully to dive briefings and directions and respect the advice of those supervising my diving activities. Recognize that additional training is recommended for participation in specialty diving activities, in other geographic areas and after periods of inactivity that exceed six months.
- 5. Adhere to the buddy system throughout every dive. Plan dives including communications, procedures for reuniting in case of separation and emergency procedures – with my buddy.
- Be proficient in dive planning (dive computer or dive table use). Make all dives no decompression dives and allow a margin of safety. Have a means to monitor depth and time underwater. Limit maximum depth to my level of training and experience. Ascend at a rate of not more than 18 metres/60 feet per minute. Be a SAFE diver – Slowly Ascend From Every dive. Make a safety stop as an added precaution, usually at 5 metres/15 feet for three minutes or longer.
- 7. Maintain proper buoyancy. Adjust weighting at the surface for neutral buoyancy with no air in my buoyancy control device. Maintain neutral buoyancy while underwater. Be buoyant for surface swimming and resting. Have weights clear for easy removal, and establish buoyancy when in distress while diving. Carry at least one surface signaling device (such as signal tube, whistle, mirror).
- 8. Breathe properly for diving. Never breath-hold or skip-breathe when breathing compressed air, and avoid excessive hyperventilation when breath-hold diving. Avoid overexertion while in and underwater and dive within my limitations.
- 9. Use a boat, float or other surface support station, whenever feasible.
- 10. Know and obey local dive laws and regulations, including fish and game and dive flag laws.

I understand the importance and purposes of these established practices. I recognize they are for my own safety and well-being, and that failure to adhere to them can place me in jeopardy when diving.

Participant's Signature	Date (Day/Month/Year)
Signature of Parent or Guardian (where applicable)	Date (Day/Month/Year)



### FLORIDA ADDENDUM NOTICE TO THE MINOR CHILD'S PARENT OR NATURAL GUARDIAN

READ THIS FORM COMPLETELY AND CAREFU	JLLY. YOU ARE
AGREEING TO LET YOUR MINOR CHILD ENGAGE	E IN A POTEN-
TIALLY DANGEROUS ACTIVITY. YOU ARE AGREEIN	
Sea Base / Scouting America , AND/	OR PADI AMERI.
CAS, INC., OR ANY OF THEIR AFFILIATE AND SUBSID	
TIONS, RESPECTIVE EMPLOYEES, OFFICERS, AGENTS	-
INDEPENDENT CONTRACTORS, DIVEPROFESSIONAL	-
AND ASSIGNS, USE REASONABLE CARE IN PROVID	
ITY, THERE IS A CHANCE YOUR CHILD MAY BE SER	
OR KILLED BY PARTICIPATING IN THIS ACTIVITY BECA	
<b>CERTAIN DANGERS INHERENT IN THE ACTIVITY WH</b>	
AVOIDED OR ELIMINATED. BY SIGNING THIS FORM	VI YOU ARE GIV-
ING UP YOUR CHILD'S RIGHT AND YOUR RIGHT TO	<b>RECOVER FROM</b>
Sea Base / Scouting America , AND/	OR PADI AMERI-
CAS, INC., OR ANY OF THEIR AFFILIATE AND SUBSID	
TIONS, RESPECTIVE EMPLOYEES, OFFICERS, AGENTS	
INDEPENDENT CONTRACTORS, DIVE PROFESSIO	
TORS AND ASSIGNS IN A LAWSUIT FOR ANY PERSO	
CLUDING DEATH, TO YOUR CHILD OR ANY PROPERT	_
RESULTS FROM THE RISKS THAT ARE A NATURAL F	
TIVITY. YOU HAVE THE RIGHT TO REFUSE TO SIGN	
	-
Sea Base / Scouting America , AND/	
ICAS, INC., OR ANY OF THEIR AFFILIATE AND SU	
PORATIONS, RESPECTIVE EMPLOYEES, OFFICERS,	<del>-</del>
TRACTORS, INDEPENDENT CONTRACTORS, DIVE F	_
INSTRUCTORS AND ASSIGNS HAVE THE RIGHT TO	
YOUR CHILD PARTICIPATE IF YOU DO NOT SIGN TH	HIS FORM.
Signature of Parent of Natural Guardian (where applicable)	Date (Day/Month/Year)
Please identify minor child/children participants below:	
(Minor Child means person under the age of 18)	

Minor Child Name

PRODUCT NO. 10348 (02/21) Version 1.0

Birth Date (Day/Month/Year)